

Paradox of Quantum Information

Subhash Kak

Department of Electrical & Computer Engineering
Louisiana State University, Baton Rouge, LA 70803, USA

February 1, 2008

Abstract

The notion of quantum information related to the two different perspectives of the global and local states is examined. There is circularity in the definition of quantum information because we can speak only of the information of systems that have been specifically prepared. In particular, we examine the final state obtained by applying unitary transformations on a single qubit that belongs to an entangled pair.

An unknown quantum state is unknowable. One may postulate it to be in a pure or mixed state although there is no way to determine the type. Interactions with the state lead to its probabilistic collapse.

One must therefore speak only of systems whose states have been specifically prepared as pure states. However, quantum gates to accomplish this are liable to have their own imprecision. The pure state has uncertainty associated with it[2].

There is circularity in the consideration of quantum information. Unlike information in the classical sense where it may be defined as characteristics of an unknown system, obtained through various measurements, we can only speak of prepared states (in principle, limited by equipment precision and measurement accuracy) and then ask questions regarding its components after certain operations have been applied on it.

Consider entangled particles[1]. Given a specific pair produced by a physical process, there is no way to characterize it accurately. For example, we wouldn't know if it is in a pure state or a mixed state, or know the nature of mixedness, if it is mixed. We must begin by preparing particles in the pure $|00\rangle$ state and then, using an appropriate 2-qubit gate, steer it into the entangled pair in a pure state.

One may view the situation of the entangled particles from the two perspectives of the system and the individual particles. In the first, the state of the two particles is considered jointly (Perspective G related to the generator of the entangled pairs). In the second perspective, one may consider each particle separately, without the knowledge that it is entangled with another particle. From this perspective (corresponding to Alice or Bob), consistency demands that operations on their qubits will provide the same results as when the particles are unentangled and in an appropriate mixed state.

Consider the entangled pure state $|\psi\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. Both particles (qubits) have equal probability of being 0 or 1. We may assume that the particles have traveled apart to A and B, so that one may operate on each separately, without influencing the other. Consider now the unitary transformation

$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

Someone who doesn't know of the entanglement of the particles may consider the particle at A to be a pure state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$. Then the application of H would rotate it to the state $|0\rangle$. The state of the two particles jointly is $\frac{1}{\sqrt{2}}(|00\rangle + |01\rangle) = |0\rangle(\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle))$. Applying H on the second particle, one obtains $|00\rangle$. This is incorrect from the Perspective G and it is the expectation based on the individual's lack of knowledge regarding the system.

From the perspective of G, we know that the entangled particles cannot individually be in pure states. The individual particles are mixed, although the entangled state is a pure state. As a mixture, the transformation H leads to the state for the pair:

$$\frac{1}{2}(|00\rangle + |01\rangle + |10\rangle - |11\rangle)$$

The H transformations on the particles available to A and B may be written as:

$$H(A) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{bmatrix}$$

$$H(B) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

Applying $H(A)$ on the entangled pair yields the state $\frac{1}{2}(|00\rangle + |01\rangle + |10\rangle - |11\rangle)$ and applying $H(B)$ on it returns the state to $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$.

The application of $H(A)$ is reversed by the application of $H(B)$. Also, $H(A)|\psi\rangle = H(B)|\psi\rangle$. The H operator applied to either of the qubits creates the same change in the state. This is surprising because of the asymmetry of the operation. It is one of the paradoxical characteristics of quantum information.

The joint state of the two particles may be represented as: $\frac{1}{2}((|0\rangle + |1\rangle)|0\rangle + (|0\rangle - |1\rangle)|1\rangle) = \frac{1}{2}(|0\rangle(|0\rangle + |1\rangle) + |1\rangle(|0\rangle - |1\rangle))$. One may view the situation to be either as the particle at A to be in the mixed state $\frac{1}{\sqrt{2}}|0\rangle + |1\rangle$ and $\frac{1}{\sqrt{2}}|0\rangle - |1\rangle$ with the particle at B to be in the mixed state $|0\rangle$ and $|1\rangle$ or the other way round, equivalently, or as a combination of the two.

Mode change From an information point of view, the nature of the state of the qubit, whether mixed or pure, is significant. When appropriate gate transformations are applied, the states of the individual qubits would transition from mixed to pure[3]. This would happen when the gate removes the entanglement between the particles. Such a transition may be measurable under certain conditions and be used to communicate information without collapsing the state function.

For example, if the output is mixed as in the case of the entangled particles, H transformation on it will leave its state unchanged and a detector after the H operator will detect 0 and 1 with equal probability. On the other hand, H transformation on the pure state (created after the entanglement has been removed) $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ will change it to $|0\rangle$. The detector now will measure 0 in each case. Given several copies of such states, one can use the H transformation to distinguish the mixed and the pure states and, therefore, detect if the gate operation was applied.

But a mode change cannot take place using a local transformation. Therefore such information transmission can be successfully accomplished only across multi-qubit gates.

Reference

1. C.H. Bennett, “Quantum information and computation,” *Phys. Today* **48** (10), 24-30 (1995).
2. S. Kak, “The initialization problem in quantum computing.” *Foundations of Physics*, **29**, 267-279 (1999); quant-ph/9805002.
3. S. Kak, quant-ph/0302118.